### **PCT**

# WORLD INTELLECTUAL PROPERTY ORGANIZATION International Bureau



## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification <sup>6</sup>:
C12Q 1/04, C12M 1/28

A1

(11) International Publication Number: WO 99/23243

(43) International Publication Date: 14 May 1999 (14.05.99)

(21) International Application Number: PCT/FI98/00854

(22) International Filing Date: 3 November 1998 (03.11.98)

(30) Priority Data: 974112 3 November 1997 (03.11.97)


(71)(72) Applicant and Inventor: HAKALEHTO, Elias [FI/FI]; Kasarmikatu 12 C 1, FIN-70110 Kuopio (FI).

(74) Agent: PITKÄNEN, Hannu; Pitkänen Oy, Savilahdentie 6 L 3, FIN-70210 Kuopio (FI). (81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, GM, HR, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

#### Published

With international search report. In English translation (filed in Finnish).

(54) Title: METHOD AND APPARATUS FOR CONCENTRATING AND SEARCHING OF MICROBIOLOGICAL SPECIMENS



#### (57) Abstract

The invention relates to a method for enriching and examining microbiological samples, in which method microbes are brought into a syringe (1) or equivalent. In a method according to the invention the enrichment culture of microbes is accomplished inside a syringe (1) or equivalent. An apparatus according to the invention comprises a syringe (1) or equivalent and into the syringe or the plunger (3) of the syringe a lead—through is manufactured in order to conduct gas or gas mixture to or from the syringe.

### FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania '	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senega!
ΑÜ	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
ΑZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana .	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav	TM	Turkmenistan
BF	Burkina Faso	GR	Greece		Republic of Macedonia	TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	П	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	¥U	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI -	Côte d'Ivoire	КР-	Democratic People's	NZ	New Zealand		
CM	Cameroon		Republic of Korea	PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

WO 99/23243 PCT/F198/00854

METHOD AND APPARATUS FOR CONCENTRATING AND SEARCHING OF MICROBIOLOGICAL SPECIMENS

The invention relates to a method for enriching and examining microbiological samples, in which method microbes are brought into a syringe or equivalent. The invention also relates to an apparatus for applying the method, the apparatus consisting of a syringe or equivalent.

Collecting microbiological sweep samples is an important part of hygiene control in industrial establishments, hospitals, laboratories and other places where the hygiene of the establishment, apparatuses and equipment is an absolute operational prerequisite. Sweep samples may also be collected from e.g. human skin or mucous membrane for clinical diagnostics.

10

15

20

30

Usually a microbiological sweep sample or a picked sample is suspended from sampling means into a buffer solution or any other appropriate solution, where it may be further examined and handled. In this case a usual method for acquiring additional information on possible microbes adhered to the sampling means is so-called subculture. This is usually accomplished by transferring the microbial suspension to be examined to a liquid culture substrate or to a solid culture substrate (e.g. a Petri dish). Thereafter the microbes in the culture substrate are incubated for at least some hours, but usually for one or more days, even weeks. During this incubation phase, the microbe to be examined is enriched to such a content that it is possible to indicate it by the indication method in use. A limiting factor for the growth of aerobic organisms during incubation may be for example oxygen. On the other hand, appropriate gases can be used in the incubation of aerobic microbes to achieve and maintain an adequately anaerobic environment. The so-called microaerophilic bacteria (e.g. Campylobacter sp.) need small oxygen contents. Also changing contents of gases may, if necessary, be conveyed to the enrichment space.

In some situations rapid completion of a microbiological analysis is crucially important e.g. for the success of a patient's treatment, in choosing cleansing measures

in hygiene control, in industrial quality control etc. Many factors have further increased the threat caused by microbes (hospital infections, new human and animal pathogenes, previously unknown industrial microbe contaminants, environmental microbiological pollution etc.). In order to be able to respond to these challenges adequately effectively, so-called rapid diagnostic methods are needed for indicating and identifying microbes.

10

15

20

25

30

The method according to international patent application PCT/FI95/00398 is aiming to collect microbiological sweep samples for further examination as easily as possible and in the case of handling pathogenic microbes as safely as possible. The above is achieved with the syringe of the invention (volume e.g. 10-50 ml) which is characterized in that the surface facing away from the plunger rod comprises an adhering substrate for microbes. The adhering substrate on the surface of the plunger may be e.g. cotton, velvet or a similar porous corresponding material that has been sterilized together with the syringe or separately (e.g. an autoclave or by radiating). Biomolecules (e.g. antibodies) that improve the adhesion of certain microbes may also, if necessary, be aseptically immobilized to the adhering substrate surface. By using the syringe and its plunger with adhering substrate surface, one avoids extra handling of microbe-containing liquids by use of a pipette, which increases safety when working with infectious microbes. The sample is transferred from the syringe onto a separate culture substrate where it is grown in a usual way. The sample may also be collected into the syringe in a usual way by sucking liquid into the syringe. In this way for example a blood sample is usually accomplished. A liquid sample may naturally be collected directly into the syringe without an injection needle or by use of a specially manufactured longer tip or a tube or similar.

The object of the present invention is to provide a method and an apparatus with which the work of further analysing microbiological samples is speeded up, which leads to fast and reliable indication, idenfication, enrichment etc.

The objects of the invention are achieved by the method and apparatus, which are characterized by what is presented in the patent claim.

The method of the invention is characterized in that the enrichment or other growth of microbes is performed inside the syringe or equivalent. This enrichment takes place in adjusted conditions and the conditions may easily be changed during the enrichment. This makes possible e.g. a safe further inoculation in a desired growth phase, because the growth in the syringe is easy to follow for example visually or using a colour indicator. The use of the syringe and its measure scale make it possible to accomplish microbiological dilution series conveniently. For example a so-called -1 dilution is accomplished by taking a ninefold volume of the dilution solution into the syringe per one sample volume. The syringe is then shaken and 9 volumes, which can be used as a sample in further analysis, are removed, and another 9 volumes of dilution solution are added, which gives a so-called -2 dilution and so on.

5

10

15

20

25

30

It is possible to change the existing conditions in the syringe for example by temperature, pH and gases conducted to the substrate. It is advantageous to use a bubbly gasflow to the substrate, in which flow the composition of the gases may be adjusted, and which also may be used to adjust the temperature and the pH. At the same time the culture is mixed, which improves the even spreading of nutrients into the substrate. It is also easy to add into the syringe, if necessary, more substrate, nutrients, selective factors and other substances in liquid form.

During incubation of aerobic microbes, the diffusion of oxygen is increased by the gas-flow conducted to the syringe, which is important in order to achieve the best possible growth speed as the availability of oxygen is usually the so-called restricting factor in aerobic microbe incubation. Apart from oxygen, also gases like nitrogen and carbon dioxide may be conducted to the substrate. With the help of these, e.g. anaerobic or microaerophilic conditions can be achieved in accordance with the type of microbes needing incubation and enrichment. The partial pressures of differant gases can be used as selective factors. Gases led to the substrate may also be used as carrier gases, which make it possible to transfer to the substrate as aerosols, in a vaporous form or by corresponding means substances that control the culture conditions or the actual cultivation or even inoculate the medium or the culture. To

WO 99/23243 4 PCT/F198/00854

accomplish this, before being transferred to the syringe, the gas may be led through a liquid or a suspension containing the component that is to be added.

From a rapid microbial detection point of view it is crucial that an adequate microbial concentration needed for reliable detection is achieved as quickly as possible. It is possible to achieve this object by using the method according to the invention e.g. for the purpose of carrying out immunological detections. Possible microbial detections, in which the method according to the invention may be used are e.g. hygiene control of interior surfaces, tools and any other surfaces, mould determination from any surfaces, hygiene control of carcases (e.g. Salmonella detection) and many other microbiological sweep sample and liquid sample detections that are required in industrial establishments, health care and environmental analytics.

5

10

15

20

25

30

When exploiting the method according to the invention in using the syringe used as sampling means for microbe enrichment, in an economical application, an appropriate selective nutrient medium to enrich the microbe may be used as growth medium immediately after sampling. Thus the enrichment may begin safely immediately after the sampling without delays caused by inoculation or transfer of the sample. This is a benefit as microbes with cells in resting state have a lag-phase before bacterial growth. E.g. the lag-phase for *Staphylococcus aureus* bacteria lasts for approximately 1,5 hours.

In hospitals and other equivalent establishments the occurance of antibiotic resistant microbial strains may be determined from interior surfaces and apparatuses, and from human skin and mucous membranes. Blood samples and any other liquid samples may be examined in a similar way. These may be patient samples or other liquid samples used in hospitals that require microbiological quality examination. During the enrichment, required antibiotics may be added to the medium. Advantages of the method are in this and many other cases besides the simple and straightforward procedure and material savings also security as the transfer of hazardous microbes in laboratories is minimized.

In an economical application of the invention, the temperature, pH and/or other conditions of the culture substrate are adjusted by gas or gas mixture that is conducted to the substrate within the syringe. The gas is conducted into the syringe in a commonly known way. As the temperature during the culture may be adjusted by the gas flow, incubation chambers or equivalent aparatuses are not necessarily needed during the culture. To heat or cool the syringe specific heating or cooling blocks e.g. Peltier elements may be used.

PCT/FI98/00854

If necessary, exhaust gases may be conducted from the enrichment space of the syringe by a tube via sterilization (e.g. a filter).

In one application of the invention the syringe is placed in a holder tip upwards during enrichment. Alternatively the syringe may be placed in a holder also tip downwards if gas is led to the enrichment space through the tip.

15

20

10

In an economical application of the invention, the growth solution and sample are transferred for further examination with the same syringe where the culture is done. Samples may be transferred in the syringe if necessary, and for this purpose the tip may be manufactured to be closed with a lid, a valve or with any other closing device. Since in enrichment or in other further treatments of samples complicated apparatuses and work phases are not needed, it may be accomplished e.g. in industrial establishments beside the production line under control. Results are gained faster as there is no need for transferring or storing the sample. Also safety risks in regard to transfer and storage are decreased.

25

30

The apparatus in accordance with the invention is characterized in, that there is a lead-through or a conduit into the syringe or into the syringe plunger for conducting gas or gas mixture into the syringe. If the rod of the plunger of the syringe is manufactured with a conduit or several conduits in the rod for conducting gas, the culture substrate may be aerated or required gases may be conducted to it in order to enhance during incubation the enrichment of aerobic microbes on one hand and anaerobic on the other hand. This gas may originate from a compressed gas bottle.

Gas may be conducted into the syringe also directly through the plunger e.g. by piercing it with an injection needle to which the gas bottle is connected by a tube or a conduit. An alternative is to conduct the gas into the syringe through a lead-through or a conduit elsewhere in the syringe. The composition of the gas may be altered during the culture e.g. in accordance with the growth phase of the microbe that is being enriched. The gas may be conducted into the syringe through a sterile filter to ensure aseptic conditions. Correspondingly exhaust gas through the tip may be filtered by use of a sterile filter. The tip of the syringe may also be closed for transportation or any other reason with a cap, a lid or a valve.

10

15

20

25

Alternatively gas may be conducted into the syringe through the tip, in which case another route must be used for exhaust gas, e.g. using a lead-through with a valve or equivalent or through the plunger or the air holes in it with the help of excess pressure. The lead-through may in this case be done using an injection needle through the plunger or the wall of the syringe.

The apparatus according to the invention comprises of a syringe with an economically transparent wall or window frame. Thus it is possible to accomplish microbial growth determination optically or visually e.g. through the wall of the syringe. Thus it is naturally possible to add e.g. indicator colour solution to the growth medium e.g. to indicate the change of pH, which indicates the change of microbial metabolism in the growth medium.

In the following the invention will be described in detail with reference to the accompanying drawings, in which

FIG.1 shows a side view of an application of the apparatus to apply the method characterized in the invention, and

FIG. 2 shows a side view of another way for conducting gas into the syringe via a lead-through.

30

The apparatus shown in FIG 1 comprises of one or more syringes 1, containing microbiological culture media, which are placed in a holder 2, tip upwards, one or

more compressed bottles 4, for storing gases or mixtures of gases, a heating/cooling jacket 5, for incoming gases, wherein the temperature of the gas is adjusted by known means. There is a lead-through in the plunger 3, of the syringe, and to it is connected a conduit 7, through which gas or gas mixture is conducted from a compressed bottle into the syringe. The conduit 7, comprises one or more control valves 11, to adjust incoming gas. Additionally the apparatus comprises of a sterile filter 8, which is placed in the incoming conduit in order to sterilize incoming gas. Exhaust gas is sterilized using another sterile filter 9, which is connected to the tip of the syringe.

- In an application as shown in FIG. 2, gas is conducted through the plunger 3, using a separate injection needle 6, or an equivalent. The injection needle comprises a conduct, such as a tube 7, or a pipe, that is connected to a compressed bottle as described above.
- In the following, the method will be described by working phases while collecting samples:

Microbes are adhered to the adhering substrate, which is either moistened (water, buffer solution) or dry and attached to the surface of the plunger 3, by sweeping the examined surface to and fro with this plunger surface. Thereafter the plunger 3, is placed in an empty syringe 1, receiver and air is pushed out through an opening in the tip of the receiver. Alternatively a sample may be collected into a syringe by sucking liquid into the syringe by normal means. The sampling should be carried out in a way that it well represents the studied object and that the sampled area is large enough.

25

20

5

Thereafter the receiver is filled by suction with a required volume of liquid (e.g. buffer solution, water or culture solution) and thereafter with some air in order to help with the mixing procedure. The apparatus is thereafter swung and shaken as a test tube to ensure microbial detachment into the liquid solution.

30

In the case that culture solution is used as a solution, the syringe may be left upside down and placed in a holder 2, in a growth temperature that is appropriate for the enrichment of the microbe intended to be indicated. In the case that gas is conducted by alternative means through the tip of the syringe, the syringe is naturally not left upside down for enrichment. The culture solution may also be added into the syringe after the microbe has first been suspended into another solution. The adjustment of temperature (or pH or any equivalent parameter) may be done by means of gas conducted through a lead-through in the plunger of the syringe. Buffered salt solution, extract solution (e.g. diluted acid or base solution), detergent solution or any equivalent solution may be used for suspending.

Suspension may be used for inoculation or it may be transferred either directly or after cultivation to the culture medium or to any other further examinations. In this case e.g. microbe identification may de done using ordinary biochemical, immunological or genetic methods. In enriching the microbe to be detected and its antigen, it is possible to exploit e.g. a method characterized in Finnish Patent 93742.

15

Although the invention is described herein with reference to applications it will be appreciated that the invention may be realized in a variety of ways within the scope of the inventive idea and the appended claims.

20


#### **CLAIMS**

15

- 1. Method for enriching and examining microbiological samples, in which method microbes are brought into a syringe (1) or equivalent, c h a r a c t e r i z e d in, that the enrichment of microbes is accomplished inside a syringe (1) or equivalent.
- Method according to claim 1, c h a r a c t e r i z e d in, that a culture substrate
   placed inside a syringe or equivalent is chosen so that it is selective to required microbes.
  - 3. Method according to claim 1 or 2, c h a r a c t e r i z e d in, that the culture substrate's temperature, pH, partial pressures of gases and/or diffusions or nutrient concentrations and/or diffusions or other conditions are controlled by means of gas or gas mixture conducted to the substrate within the syringe.
  - 4. Method according to one or several of the claims 1 to 3, c h a r a c t e r i z e d in, that a sample containing microbes is collected into the syringe prior to addition of growth medium.
  - 5. Method according to one or several of the claims 1 to 4, c h a r a c t e r i z e d in, that a syringe is placed tip upwards in a holder (2) for the enrichment period.
- 6. Method according to one or several of the claims 1 to 4, c h a r a c t e r i z e d in, that a syringe is placed tip downwards in a holder for the enrichment period and in that gas is conducted into the syringe through the tip.
- 7. Method according to one or several of the claims 1 to 6, c h a r a c t e r i z e d in,
  that a culture solution is transferred for further examination using the same syringe where the culture was accomplished.

- 8. Method according to one or several of the claims 1 to 7, c h a r a c t e r i z e d in, that dilutions according to microbiological dilution series may be accomplished in the syringe prior to further inoculations.
- 9. Apparatus for applying the method according to one or several of the claims 1 to 8, the apparatus comprising a syringe (1) or equivalent, c h a r a c t e r i z e d in, that a lead-through is manufactured in the syringe or the plunger (3) of the syringe, in order to conduct gas or gas mixture in or out of the syringe.
- 10. Apparatus according to claim 9, c h a r a c t e r i z e d in, that the apparatus comprises one or more compressed bottles (4) in order to conduct gas or gas mixture from the bottles into the syringe.
- 11. Apparatus according to claims 9 and/or 10, c h a r a c t e r i z e d in, that the apparatus comprises a heating/cooling jacket (5), in which the temperature of gas is adjusted as required prior to conducting it to the syringe.
- 12. Apparatus according to claim 9, c h a r a c t e r i z e d in, that into a syringe or a plunger (3) of the syringe an injection needle (6) or equivalent is used to produce a20 lead-through.
  - 13. Apparatus according to claim 12, c h a r a c t e r i z e d in, that a tube (7) or a pipe is attached to the injection needle in order to conduct gas in or out of the syringe.
- 25 14. Apparatus according to one or several of the claims 9 to 13, c h a r a c t e r i z e d in, that the apparatus comprises a lid, cap or a valve to close the tip of the syringe.
  - 15. Apparatus according to one or several of the claims 9 to 13, c h a r a c t e r i z e d in, that the apparatus comprises a transparent wall or a window frame in order to monitor the growth of the microbial culture.

16. Apparatus according to one or several of the claims 9 to 13, c h a r a c t e r ı z e d in, that the apparatus comprises one or more sterile filters (8,9) in order to conduct incoming and/or exhaust gas or gas mixture through a sterile filter.



## INTERNATIONAL SEARCH REPORT

International application No.

PCT/FI 98/00854

		101711 3070	330 (
A. CLASSIFICATION	N OF SUBJECT MATTER		
IPC6: C120 1/04	, C12M 1/28 I Patent Classification (IPC) or to both nati		
		onal classification and IPC	
B. FIELDS SEARCH	Searched (classification system followed by	classification symbols)	
		Casamonion of moony	
IPC6: C12Q, C12	other than minimum documentation to the	weent that such documents are included	in the fields searched
		extent that such documents are included	m the rend searches
SE,DK,FI,NO cla			
Electronic data base cons	ulted during the international search (name o	of data base and, where practicable, sear	ch terms used)
WPI, EPODOC			
C. DOCUMENTS C	ONSIDERED TO BE RELEVANT		
Category* Citation of	document, with indication, where appr	opriate, of the relevant passages	Relevant to claim No.
	177 C (STIG STÄLHANDSKE), 1 5.04.96)	15 April 1996	1,2,4-8
	·		
	4491 A (SVEN E. LINDGREN), 8.10.77)	18 October 1977	1,2,4-8
	440.54		
		,	·
	,		
Further document	nts are listed in the continuation of Box	C. X See patent family ann	ex.
Special categories of		"T" later document published after the i	nternational filing date or priority
"A" document defining the	e general state of the art which is not considered evance	date and not in conflict with the ap the principle or theory underlying the	
"E" erlier document but p	oublished on or after the international filing date throw doubts on priority claim(s) or which is	"X" document of particular relevance: the considered novel or cannot be consisted when the document is taken along the consisted when the document is taken along the consistency of the	dered to involve an inventive
cited to establish the special reason (as spe	publication date of another citation or other cified)	"Y" document of particular relevance: ti	he claimed invention cannot be
means	o an oral disclosure, use, exhibition or other	considered to involve an inventive a combined with one or more other s being obvious to a person skilled in	uch documents, such combination
"P" document published the priority date claim	prior to the international filing date but later than- med	"&" document member of the same pate	
Date of the actual co	mpletion of the international search	Date of mailing of the internationa 2 6 -0	l search report 2- 1999
22 February 19	999	Authorized officer	·
Name and mailing ad Swedish Patent Of		Authorized officer	
Box 5055, S-102 42	STOCKHOLM	Hampus Rystedt	
Facsimile No. +46	8 666 02 86	Telephone No. + 46 8 782 25 00	J

## INTERNATIONAL SEARCH REPORT

Information on patent family members

02/02/99

International application No. PCT/FI 98/00854

Patent document cited in search report		Publication date		Patent family member(s)	Publication date
SE	503177 C	15/04/96	AU	2312697 A	20/10/98
	•		SE	9500682 A	15/04/96
			WO	9842821 A	01/10/98
			SE	9700378 D	00/00/00
US	4054491 A	18/10/77	DE	2453858 A	15/05/75
			DK	136430 B,C	10/10/77
			DK	591174 A	14/07/75
			FI	52739 B,C	01/08/77
			FI	326574 A	14/05/75
			FR	2250998 A,B	06/06/75
			GB	1489229 A	19/10/77
	•		JP	932068 C	07/11/78
			JP	50077583 A	24/06/75
			JP	53008791 B	31/03/78
			NL	161503 B,C	17/09/79
			NL	7414821 A	15/05/75
			SE	377811 B,C	28/07/75
			SE	7315344 A	14/05/75